

Yukawa Institute Kyoto

YITP-01-55
hep-th/0108155
August 2001

On the integrability of classical Ruijsenaars-Schneider Model of BC_2 type

V. I. Inozemtsev¹ and R. Sasaki

Yukawa Institute for Theoretical Physics
Kyoto University, Kyoto 606-8502, Japan

Abstract

The problem of finding most general form of the classical integrable relativistic models of many-body interaction of the BC_n type is considered. In the simplest nontrivial case of $n=2$, the extra integral of motion is presented in explicit form within the ansatz similar to the nonrelativistic Calogero-Moser models. The resulting Hamiltonian has been found by solving the set of two functional equations.

¹ permanent address: BLTP JINR, 141980 Dubna, Moscow Region, Russia

The one-parameter extensions of the Calogero-Moser integrable particle systems first proposed by Ruijsenaars and Schneider [1,2] have been extensively studied during last decade [3,4,6-17]. It turned out that both classical and quantum versions of these systems (often called “relativistic” but, as explained in [6], this term is misleading) were connected with various branches of mathematical physics. The quantum RS models with trigonometric type of interaction have been found to be relevant in the theory of Macdonald polynomials and the integrability of their elliptic version has been investigated within the Yang-Baxter approach [8,9]. The applications of classical models range from characteristics of multisoliton solutions of the sine-Gordon equation to the Seiberg-Witten theory [13].

The original n -particle RS model has the Hamiltonian

$$H_n = \sum_{j=1}^n \text{ch} p_j \prod_{k \neq j}^n F(q_j - q_k),$$

where p, q are canonical variables with Poisson brackets $\{p_j, p_k\}_P = \{q_j, q_k\}_P = 0$, $\{q_j, p_k\}_P = \delta_{jk}$. It is translation invariant and can be associated with A_{n-1} root system. Extensions to other root systems were proposed [7,10-17] in ways similar to the Olshanetsky-Perelomov construction for Calogero-Moser models [18]. One can find the first of them in the paper by Schneider [4] who considered the non-translation-invariant model with an external field and the Hamiltonian

$$H_S = H_n + \sum_{j=1}^n W(q_j).$$

The Schneider model has been found to be integrable for hyperbolic choice of the two-particle interaction,

$$F^2(q) = 1 + g^2 \text{sh}^2(aq), \quad W(q) = A^2 \exp(\pm 2aq),$$

which coincides upon the limit

$$p_j \rightarrow p_j/c, \quad q_j \rightarrow q_j c, \quad g \rightarrow g/c, \quad A \rightarrow A/c, \quad c \rightarrow \infty$$

with well-known Adler system [5]. The next step has been done in the papers by van Diejen [7] who has studied the quantum RS systems and has found their nine-parametric extension (with one constraint) of the BC_n type in the most general elliptic case. The

results of [7] have been proved rigorously for $n = 2$ only. Komori and Hikami [10,11] also considered the elliptic potentials starting from the paper by Hasegawa [9] within the framework of quantum inverse scattering method and gave a rigorous proof of the commutativity of van Diejen operators for arbitrary n and with no constraint among couplings. As for classical models, the Lax operators were obtained in explicit form only for very particular case of Z_2 folding of the original A_{2n} and A_{2n+1} RS model [14-16].

The aim of this Letter is to find most general form of classical integrable extensions of the RS hamiltonian of the BC -type. The general procedure of constructing the Lax pairs for these systems is still lacking, so we shall consider the simplest but nontrivial systems of two degrees of freedom. In this two-particle case, one needs to find only one extra integral of motion $J(p_1, p_2, q_1, q_2)$ so as to prove the integrability of the system.

We shall consider the following Ansatz for the two-particle Hamiltonian and an extra integral of motion J ,

$$H_2 = V_1(q_1, q_2) \operatorname{ch} p_1 + V_2(q_1, q_2) \operatorname{ch} p_2 + W(q_1, q_2), \quad (1)$$

$$J = \lambda(q_1, q_2) \operatorname{ch} p_1 \operatorname{ch} p_2 + \mu(q_1, q_2) \operatorname{sh} p_1 \operatorname{sh} p_2 + \rho_1(q_1, q_2) \operatorname{ch} p_1 + \rho_2(q_1, q_2) \operatorname{ch} p_2 + \tau(q_1, q_2), \quad (2)$$

where $V_{1,2}$, W , λ , μ , $\rho_{1,2}$ and τ are yet unspecified functions of particle coordinates. We shall suppose that H_2 and J are symmetric with respect to the permutation of particles,

$$V_2(q_1, q_2) = V_1(q_2, q_1), \quad W(q_1, q_2) = W(q_2, q_1), \quad (3)$$

$$\lambda(q_1, q_2) = \lambda(q_2, q_1), \quad \mu(q_1, q_2) = \mu(q_2, q_1), \quad \rho_2(q_1, q_2) = \rho_1(q_2, q_1), \quad \tau(q_1, q_2) = \tau(q_2, q_1). \quad (4)$$

The calculation of the Poisson bracket $\{H_2, J\}_P$ and equating it to zero results in an overdetermined system of 10 equations for 8 unknown functions,

$$V_1 \frac{\partial \lambda}{\partial q_1} - \lambda \frac{\partial V_1}{\partial q_1} - \mu \frac{\partial V_1}{\partial q_2} = 0, \quad V_2 \frac{\partial \lambda}{\partial q_2} - \mu \frac{\partial V_2}{\partial q_1} - \lambda \frac{\partial V_2}{\partial q_2} = 0, \quad (5)$$

$$V_1 \frac{\partial \mu}{\partial q_1} - \mu \frac{\partial V_1}{\partial q_1} - \lambda \frac{\partial V_1}{\partial q_2} = 0, \quad V_2 \frac{\partial \mu}{\partial q_2} - \lambda \frac{\partial V_2}{\partial q_1} - \mu \frac{\partial V_2}{\partial q_2} = 0, \quad (6)$$

$$V_1 \frac{\partial \rho_1}{\partial q_1} - \rho_1 \frac{\partial V_1}{\partial q_1} = 0, \quad V_2 \frac{\partial \rho_2}{\partial q_2} - \rho_2 \frac{\partial V_2}{\partial q_2} = 0, \quad (7)$$

$$V_1 \frac{\partial \tau}{\partial q_1} - V_2 \frac{\partial \mu}{\partial q_2} - \rho_1 \frac{\partial W}{\partial q_1} = 0, \quad V_2 \frac{\partial \tau}{\partial q_2} - V_1 \frac{\partial \mu}{\partial q_1} - \rho_2 \frac{\partial W}{\partial q_2} = 0, \quad (8)$$

$$V_1 \frac{\partial \rho_2}{\partial q_1} - \rho_1 \frac{\partial V_2}{\partial q_1} - \mu \frac{\partial W}{\partial q_2} - \lambda \frac{\partial W}{\partial q_1} = 0, \quad V_2 \frac{\partial \rho_1}{\partial q_2} - \rho_2 \frac{\partial V_1}{\partial q_2} - \mu \frac{\partial W}{\partial q_1} - \lambda \frac{\partial W}{\partial q_2} = 0. \quad (9)$$

To solve the system (5-9), it is useful to start from four equations (5,6). They can be easily linearized and the general solution which takes into account also the symmetry (3) reads

$$V_1 = F(q_1 - q_2)G(q_1 + q_2)R(q_1), \quad V_2 = F(q_1 - q_2)G(q_1 + q_2)R(q_2), \quad (10)$$

where F, G, R are (still unknown) functions of one argument, and

$$\lambda = -\frac{1}{2}[F^2(q_1 - q_2) + G^2(q_1 + q_2)]R(q_1)R(q_2), \quad \mu = \frac{1}{2}[F^2(q_1 - q_2) - G^2(q_1 + q_2)]R(q_1)R(q_2). \quad (11)$$

The solution of (7) has the form

$$\rho_1 = V_1 d(q_2), \quad \rho_2 = V_2 d(q_1). \quad (12)$$

The equations (8) can be rewritten as

$$\frac{\partial \tau}{\partial q_1} = \frac{R(q_2)}{R(q_1)} \frac{\partial \mu}{\partial q_2} + d(q_2) \frac{\partial W}{\partial q_1}, \quad \frac{\partial \tau}{\partial q_2} = \frac{R(q_1)}{R(q_2)} \frac{\partial \mu}{\partial q_1} + d(q_1) \frac{\partial W}{\partial q_2}. \quad (13)$$

These equations with the use of (11) can be cast into the form

$$\frac{\partial \tau}{\partial q_1} = R(q_2) \left[\frac{1}{2} R'(q_2) (F^2 - G^2) - R(q_2) (FF' + GG') \right] + d(q_2) \frac{\partial W}{\partial q_1}, \quad (14)$$

$$\frac{\partial \tau}{\partial q_2} = R(q_1) \left[\frac{1}{2} R'(q_1) (F^2 - G^2) + R(q_1) (FF' - GG') \right] + d(q_1) \frac{\partial W}{\partial q_2}, \quad (15)$$

where prime means differentiation of the function with respect to its argument. Now the equality of second mixed derivatives of τ obtained from (14,15) yields

$$\begin{aligned} & \frac{\partial^2 W}{\partial q_1 \partial q_2} (d(q_2) - d(q_1)) + d'(q_2) \frac{\partial W}{\partial q_1} - d'(q_1) \frac{\partial W}{\partial q_2} \\ & + \frac{\partial}{\partial q_2} \left[\frac{1}{2} R(q_2) R'(q_2) (F^2 - G^2) - R^2(q_2) (FF' + GG') \right] \\ & - \frac{\partial}{\partial q_1} \left[\frac{1}{2} R(q_1) R'(q_1) (F^2 - G^2) + R^2(q_1) (FF' - GG') \right] = 0. \end{aligned} \quad (16)$$

It is useful to introduce instead of F, G and R new functions by

$$F = (1 + f)^{1/2}, \quad G = (1 + g)^{1/2}, \quad R = (1 + r)^{1/2}. \quad (17)$$

The substitution of (10-12, 17) into the last pair of equations (9) yields after some transformations

$$\frac{\partial W}{\partial q_1} = \frac{1}{2} [(f' + g')(d(q_2) - d(q_1)) - (1 + f)(d'(q_1) + d'(q_2)) + (1 + g)(d'(q_2) - d'(q_1))], \quad (18)$$

$$\frac{\partial W}{\partial q_2} = \frac{1}{2}[(f' - g')(d(q_2) - d(q_1)) - (1 + f)(d'(q_1) + d'(q_2)) - (1 + g)(d'(q_2) - d'(q_1))]. \quad (19)$$

The compatibility condition for the equations (18-19) reads

$$2(f''(q_1 - q_2) - g''(q_1 + q_2))(d(q_2) - d(q_1)) + (f(q_1 - q_2) - g(q_1 + q_2))(d''(q_2) - d''(q_1)) - 3f'(q_1 - q_2)(d'(q_1) + d'(q_2)) + 3g'(q_1 + q_2)(d'(q_1) - d'(q_2)) = 0. \quad (20)$$

The equation (16) can be transformed with the use of (17-19) to the form

$$\begin{aligned} & 2(f''(q_1 - q_2) - g''(q_1 + q_2))(r(q_2) - r(q_1)) + (f(q_1 - q_2) - g(q_1 + q_2))(r''(q_2) - r''(q_1)) \\ & - 3f'(q_1 - q_2)(r'(q_1) + r'(q_2)) + 3g'(q_1 + q_2)(r'(q_1) - r'(q_2)) \\ = & (d(q_1) - d(q_2))[(d''(q_1) + d''(q_2))(g(q_1 + q_2) - f(q_1 - q_2)) + 3f'(q_1 - q_2)(d'(q_2) - d'(q_1)) \\ & + 3g'(q_1 + q_2)(d'(q_1) + d'(q_2))] + 2(g(q_1 + q_2) - f(q_1 - q_2))(d'^2(q_1) - d'^2(q_2)). \end{aligned} \quad (21)$$

One is left with two functional equations (20, 21). The first of them allows one to find the functions f, g and d while the second one can be regarded as inhomogeneous functional equation for the function $r(q)$. The most general solution to (20) with $f = g$ has been found in [19],

$$f(q) = g(q) = f_0 \wp(q), \quad (22)$$

where $\wp(q)$ is the Weierstrass function with two arbitrary periods $2\omega_1, 2\omega_2$,

$$d(q) = \sum_{i=0}^3 \alpha_i \wp(q + \omega_i), \quad (23)$$

where we use the notation $\omega_0 = 0$, $\omega_3 = -\omega_1 - \omega_2$, and $\{\alpha_i\}$ are four arbitrary real parameters.

This solution has to be used in equation (21) which is bilinear in $d(q)$. Plugging (22-23) into right-hand side of (21) results, besides the terms quadratic in $\{\alpha_i\}$, in 6 terms proportional to $\alpha_i \alpha_k$ with $i \neq k$. However, it is possible with the use of *Mathematica* to show that all these cross-terms vanish and the right-hand side of (21) acquires simple form

$$8 \sum_{i=0}^3 \alpha_i^2 \wp'(q_1 + \omega_i) \wp'(q_2 + \omega_i) [\wp(q_2 + \omega_i) - \wp(q_1 + \omega_i)].$$

It allows one to find the general solution to (21),

$$r(q) = \sum_{i=0}^3 [r_i \wp(q + \omega_i) + \alpha_i^2 \wp^2(q + \omega_i)], \quad (24)$$

where $\{r_i\}$ is yet another set of four arbitrary real parameters. After solving equations (18-19) the resulting Hamiltonian reads

$$H_2 = \sum_{j=1}^2 \text{ch} p_j \prod_{k \neq j} \{[1 + f(q_j - q_k)][1 + f(q_j + q_k)][1 + r(q_j)]\}^{1/2} + W(q_1, q_2), \quad (25)$$

where f, g and r are given by (22,24) and

$$W(q_1, q_2) = - \sum_{i=0}^3 [f_0 \alpha_i \wp(q_1 + \omega_i) \wp(q_2 + \omega_i)] - d(q_1) - d(q_2). \quad (26)$$

In the papers [7,10,11] the Hamiltonian was written in terms of the products of the Weierstrass sigma functions. Our solution (24) for $1 + r(q)$ can be also cast into this form by using the formulas

$$\begin{aligned} \wp(q) \wp(q + \omega_i) &= \wp(\omega_i) [\wp(q) + \wp(q + \omega_i)] - \wp^2(\omega_i) + \wp''(\omega_i)/2, \\ \wp(q) - \wp(\alpha) &= \frac{\sigma(\alpha - q) \sigma(\alpha + q)}{\sigma^2(q) \sigma^2(\alpha)}. \end{aligned}$$

The second invariant J is determined by the formulas (11-12) and (14-15). The explicit expression for the function $\tau(q_1, q_2)$ can be also obtained by the integration of (14-15). In the elliptic case, it reads

$$\begin{aligned} \tau(q_1, q_2) &= -\frac{1}{2} [f(q_1 - q_2) + f(q_1 + q_2)] - d(q_1) d(q_2) \\ &\quad + f_0 \sum_{i=0}^3 [r_i \psi(q_1 + \omega_i, q_2 + \omega_i) + \alpha_i^2 \chi(q_1 + \omega_i, q_2 + \omega_i)] \\ &\quad - f_0 \sum_{i>k=0} \frac{\alpha_i \alpha_k}{\wp(\omega_i - \omega_k)} \prod_{\gamma=1}^2 \wp(q_\gamma + \omega_i) \wp(q_\gamma + \omega_k), \end{aligned} \quad (27)$$

where $f(q)$ and $d(q)$ are given by (22-23) and

$$\begin{aligned} \psi(q_1, q_2) &= \wp(q_1) \wp(q_2) - \frac{\wp(q_1) \wp'^2(q_2) + \wp(q_2) \wp'^2(q_1)}{4(\wp(q_1) - \wp(q_2))^2}, \\ \chi(q_1, q_2) &= -\frac{\wp^2(q_1) \wp'^2(q_2) + \wp^2(q_2) \wp'^2(q_1)}{4(\wp(q_1) - \wp(q_2))^2}. \end{aligned}$$

Taking correct hyperbolic (trigonometric) and rational limits of (22-26) is a nontrivial task. To do this, we have solved again the equations (21-22) with the use of *Mathematica*. The results are given by (25) with the following sets of functions f, g, d, r, W and τ :

(a) Hyperbolic case:

$$f(q) = g(q) = \frac{f_0}{\text{sh}^2 aq}, \quad (28)$$

$$d(q) = \frac{\alpha_0}{\text{sh}^2 aq} + \frac{\alpha_1}{\text{ch}^2 aq} + \alpha_2 \text{ch} 2aq + \alpha_3 \text{ch} 4aq, \quad (29)$$

$$r(q) = \frac{r_0}{\text{sh}^2 aq} + \frac{r_1}{\text{ch}^2 aq} + r_2 \text{ch} 2aq + r_3 \text{ch} 4aq \\ + \frac{\alpha_0^2}{\text{sh}^4 aq} + \frac{\alpha_1^2}{\text{ch}^4 aq} + \alpha_2 \alpha_3 \text{ch} 6aq + \frac{\alpha_3^2}{2} \text{ch} 8aq, \quad (30)$$

$$W(q_1, q_2) = - \sum_{i=1}^2 d(q_i) - f_0 \left(\frac{\alpha_0}{\text{sh}^2 aq_1 \text{sh}^2 aq_2} - \frac{\alpha_1}{\text{ch}^2 aq_1 \text{ch}^2 aq_2} + 4\alpha_3 \text{ch} 2aq_1 \text{ch} 2aq_2 \right), \quad (31)$$

$$\begin{aligned} \tau(q_1, q_2) = & -d(q_1)d(q_2) - (\text{sha}(q_1 - q_2) \text{sha}(q_1 + q_2))^{-2} f_0 [(\text{ch} 2aq_1 \text{ch} 2aq_2 - 1)/2 \\ & + r_0(\text{ch}^2 aq_1 + \text{ch}^2 aq_2) + r_1(\text{sh}^2 aq_1 + \text{sh}^2 aq_2) + r_2(\text{sh}^2 aq_1 \text{ch} 2aq_1 \text{ch}^2 aq_2 \\ & + \text{ch}^2 aq_1 \text{sh}^2 aq_2 \text{ch} 2aq_2) + r_3(\text{sh}^2 2aq_1 \text{sh}^2 2aq_2 + \text{sh}^2 aq_1 \text{ch}^2 aq_2 + \text{sh}^2 aq_2 \text{ch}^2 aq_1) \\ & + \alpha_0^2(\coth^2 aq_1 + \coth^2 aq_2) + \alpha_1^2(\tanh^2 aq_1 + \tanh^2 aq_2) + \alpha_2 \alpha_3(\text{ch} 2aq_1 + \text{ch} 2aq_2) \\ & \times (4 \text{ch} 2aq_1 \text{ch} 2aq_2 - 3)(\text{ch} 2aq_1 \text{ch} 2aq_2 - 1)/4 + \alpha_3^2(8 \text{ch}^3 2aq_1 \text{ch}^3 2aq_2 \\ & + 9 \text{ch} 2aq_1 \text{ch} 2aq_2 - 4 \text{ch} 2aq_1 \text{ch} 2aq_2(\text{ch} 2aq_1 + \text{ch} 2aq_2)^2 - 1)/4] \\ & - f_0 [16\alpha_0 \alpha_1(\text{sh} 2aq_1 \text{sh} 2aq_2)^{-2} + \alpha_0 \alpha_2(\text{sha} q_1 \text{sha} q_2)^{-2}(1 + 2 \text{sh}^2 aq_1 + 2 \text{sh}^2 aq_2) \\ & + \alpha_1 \alpha_2(\text{cha} q_1 \text{cha} q_2)^{-2}(1 - 2 \text{ch}^2 aq_1 - 2 \text{ch}^2 aq_2) + \alpha_0 \alpha_3(16(\text{sh}^2 aq_1 + \text{sh}^2 aq_2) \\ & + 8(\text{ch}^2 aq_1 \text{sh}^{-2} aq_2 + \text{ch}^2 aq_2 \text{sh}^{-2} aq_1) + (\text{sha} q_1 \text{sha} q_2)^{-2}) + \alpha_1 \alpha_3(16(\text{ch}^2 aq_1 \\ & + \text{ch}^2 aq_2) - 8(\text{sh}^2 aq_1 \text{ch}^{-2} aq_2 + \text{sh}^2 aq_2 \text{ch}^{-2} aq_1) - (\text{cha} q_1 \text{cha} q_2)^{-2})], \quad (32) \end{aligned}$$

where $\{r_i, \alpha_i\}$ is the set of 8 arbitrary real parameters which does not coincide with those introduced for general elliptic case in (23),(24) (the same remark holds also for subsequent rational and asymmetric hyperbolic models).

(b) Rational case:

$$f(q) = g(q) = f_0 q^{-2}, \quad (33)$$

$$d(q) = \alpha_0 q^{-2} + \alpha_1 q^2 + \alpha_2 q^4 + \alpha_3 q^6, \quad (34)$$

$$r(q) = r_0 q^{-2} + r_1 q^2 + r_2 q^4 + r_3 q^6 \\ + \alpha_0^2 q^{-4} + (2\alpha_1 \alpha_3 + \alpha_2^2) q^8 + 2\alpha_2 \alpha_3 q^{10} + \alpha_3^2 q^{12}, \quad (35)$$

$$W(q_1, q_2) = - \sum_{i=1}^2 d(q_i) - f_0 [\alpha_0 (q_1 q_2)^{-2} + \alpha_2 (q_1^2 + q_2^2) + \alpha_3 (q_1^4 + q_2^4 + 3q_1^2 q_2^2)], \quad (36)$$

$$\begin{aligned} \tau(q_1, q_2) = & -\frac{1}{2} [f(q_1 - q_2) + f(q_1 + q_2)] - d(q_1)d(q_2) \\ & - \frac{f_0}{(q_1^2 - q_2^2)^2} \left[2r_0 + 2r_1 q_1^2 q_2^2 + r_2 q_1^2 q_2^2 + 2r_3 q_1^4 q_2^4 + (q_1^2 + q_2^2)(\alpha_0^2 q_1^{-2} q_2^{-2} + \alpha_2^2 q_1^4 q_2^4 \right. \\ & \quad \left. + \alpha_3^2 q_1^6 q_2^6 + \alpha_2 \alpha_3 q_1^4 q_2^4 (q_1^2 + q_2^2)) + \alpha_1 \alpha_3 q_1^2 q_2^2 (q_1^6 + q_2^6 + q_1^2 q_2^2 (q_1^2 + q_2^2)) \right] \\ & - f_0 \left[\alpha_0 \alpha_1 (q_1^{-2} + q_2^{-2}) + \alpha_0 \alpha_2 (q_1^2 q_2^{-2} + q_2^2 q_1^{-2}) \right. \\ & \quad \left. + \alpha_0 \alpha_3 (q_1^4 q_2^{-2} + q_2^4 q_1^{-2} + 3(q_1^2 + q_2^2)) + \alpha_1 \alpha_2 q_1^2 q_2^2 \right], \end{aligned} \quad (37)$$

where $f(q)$ and $d(q)$ are defined by (33-34) and $\{\alpha_i, r_i\}$ is the set of 8 real parameters which do not coincide with those in (27). There is yet another hyperbolic degeneration which corresponds to the generalization of Schneider case [4] of interaction with external field,

$$f(q) = \frac{f_0}{\text{sh}^2 a q}, \quad g = 0, \quad (38)$$

$$d(q) = \alpha_0 e^{-2aq} + \alpha_1 e^{2aq} + \alpha_2 e^{-4aq} + \alpha_3 e^{4aq}, \quad (39)$$

$$\begin{aligned} r(q) = & r_0 e^{-2aq} + r_1 e^{2aq} + r_2 e^{-4aq} + r_3 e^{4aq} \\ & + 2\alpha_0 \alpha_2 e^{-6aq} + 2\alpha_1 \alpha_3 e^{6aq} + \alpha_2^2 e^{-8aq} + \alpha_3^2 e^{8aq}, \end{aligned} \quad (40)$$

$$W(q_1, q_2) = - \sum_{i=1}^2 d(q_i) - 2f_0 \left(\alpha_2 e^{-2a(q_1+q_2)} + \alpha_3 e^{2a(q_1+q_2)} \right), \quad (41)$$

$$\begin{aligned} \tau(q_1, q_2) = & -d(q_1)d(q_2) - f_0 [(2 \text{sh}^2 a(q_1 - q_2))^{-1} [1 + r_0 (e^{-2aq_1} + e^{-2aq_2})/2 \\ & + r_1 (e^{2aq_1} + e^{2aq_2})/2 + r_2 e^{-2a(q_1+q_2)} + r_3 e^{2a(q_1+q_2)} + \alpha_0 \alpha_2 e^{-2a(q_1+q_2)} (e^{-2aq_1} + e^{-2aq_2}) \\ & + \alpha_1 \alpha_3 e^{2a(q_1+q_2)} (e^{2aq_1} + e^{2aq_2}) + \alpha_2^2 e^{-4a(q_1+q_2)} + \alpha_3^2 e^{4a(q_1+q_2)}] \\ & + 2\alpha_0 \alpha_3 (e^{2aq_1} + e^{2aq_2}) + 2\alpha_1 \alpha_2 (e^{-2aq_1} + e^{-2aq_2}) + 4\alpha_2 \alpha_3 \text{ch} 2a(q_1 - q_2)]. \end{aligned} \quad (42)$$

The original Schneider's result [4] for two particles corresponds to the case of all $r_i=0$, $\alpha_2 = \alpha_3 = 0$ in (38-41).

Let us summarize our results. We have demonstrated the Liouville integrability of the classical Ruijsenaars-Schneider Hamiltonian of the BC_2 type. It contains nine arbitrary coupling constants with no constraints in the general elliptic case and its hyperbolic and rational degenerations. Its form (25) is suitable for taking "nonrelativistic" limit $p \rightarrow p/c$, $q \rightarrow cq$, $c \rightarrow \infty$: it is necessary to rescale $f_0 \rightarrow f_0/c^2$, $r_i \rightarrow r_i/c^2$, $\alpha_i \rightarrow 0$ for getting most

general five-parametric two-particle Calogero-Moser Hamiltonian given in [19]. It would be of interest to construct the Lax representation for our general nine-parametric case so as to prove the classical integrability for arbitrary number of particles. At present time it was done only for very particular cases [14-17].

Acknowledgments. V.I.I. is supported by JSPS long term fellowship. R.S. is partially supported by the Grant-in aid from the Ministry of Education, Culture, Sports, Science and Technology, Japan, priority area (#707) “Supersymmetry and unified theory of elementary particles”.

References

1. S.N.M. Ruijsenaars and H. Schneider. Ann. Phys. (NY) **170** (1986) 370.
2. S.N.M. Ruijsenaars. Commun. Math. Phys. **110** (1987) 191; **115** (1988) 127.
3. M. Bruschi and F. Calogero. Commun. Math. Phys. **109** (1987) 481.
4. H. Schneider. Physica **D26**, (1987) 203.
5. M. Adler. Commun. Math. Phys. **55** (1977) 195.
6. H.W. Braden and R. Sasaki. Prog. Theor. Phys. **97** (1997) 1003.
7. J.F. van Diejen. J. Math. Phys. **35** (1994) 2983; **36** (1995) 1299.
8. A. Antonov, K. Hasegawa and A. Zabrodin. Nucl. Phys. **B503** (1997) 747.
9. K. Hasegawa. Commun. Math. Phys. **187** (1997) 289.
10. Y. Komori and K. Hikami. J. Phys. **A30** (1997) 434.
11. Y. Komori and K. Hikami. J. Math. Phys. **39** (1998) 6175.
12. K. Hasegawa, T. Ikeda and T. Kikuchi. J. Math. Phys. **40** (1999) 4549.
13. H.W. Braden, A. Marshakov, A. Mironov, A. Morozov. Nucl. Phys. **B558** (1999) 371.
14. B. Hou, W.-L. Yang. J. Math. Phys. **41** (2000) 357.

- 15. K. Chen, B. Hou and W.-L. Yang. J. Math. Phys. **41** (2000) 8132.
- 16. K. Chen and B. Hou. The D_n Ruijsenaars-Schneider Model, hep-th/0102036.
- 17. J. Avan and G. Rollet. Structures in BC_N Ruijsenaars-Schneider Models, nlin.SI/0106015.
- 18. M.A. Olshanetsky and A.M. Perelomov. Phys. Reports **71** (1981) 313.
- 19. V.I. Inozemtsev. J. Phys. **A17** (1984) 815.